

## **Light and temperature interactions in promoting lettuce seed germination**

Samuel Contreras<sup>1</sup>, David Tay<sup>2</sup> and Mark Bennett<sup>1</sup>

<sup>1</sup>Department of Horticulture and Crop Science, Ohio State University, Columbus, OH, 43210, USA.

<sup>2</sup>Ornamental Plant Germplasm Center, Ohio State University, Columbus, OH, 43210, USA.

Lettuce seeds were used with the objective of determining the effect of temperature, light and their interactions in promoting germination. At 20°C and constant light, the seed presented 100% germination. Different treatments evaluated germination under dark conditions, with or without a light break (LB) 48 h after sowing, and with different combination of temperatures pre (soaking temperature, ST) and post (germination temperature, GT) the LB. Germination at constant 20°C without LB was less than 5%, and with LB it was around 30%. However, germination was close to 100% at GT of 20°C when LB was applied after a ST of 10°C, and around 50% under the same conditions but without LB. When GT was 30°C and LB was applied, germination was less than 3% with ST= 30°C, less than 10% with ST=20°C, and around 100% when ST=10°C. With ST and GT of 10°C and 30°C, respectively, and no LB, germination was less than 5%. Germination at 10°C constant, with and without LB was around 90% and 0% respectively. When ST was 40°C and LB was applied, germination was around 40% at GT= 20°C, but less than 3% with GT= 30°C. In summary, a severe inhibition of germination was observed when seeds were germinated in dark, which was partially reversed by either a light treatment or soaking at 10°C, and fully reversed when both treatments were applied together. Inhibition of lettuce germination at 30°C was observed when this temperature was applied after a light treatment, but not when applied before. Possible implications of these germination results for the phytochrome mechanism of action are discussed.